



ANTHELMINTIC AND ANTIMICROBIAL PROPERTIES OF LEAVES OF *BRUGUIERA GYMNORRHIZA (L.) LAM.*

***K. R. Vandana, K. Bhagya Deepthi, K. Yugandhar, P.S. Kiran, A. Saravanakumar**

*Sree Vidyankethan College of Pharmacy, Sree Sainath nagar,
Tirupati PIN: 517-102, Andhra Pradesh.

ABSTRACT

The objective of the present investigation was to determine the anthelmintic and antimicrobial activity of petroleum ether extract of the leaves of *Bruguiera gymnorrhiza (L.) Lam.* Anthelmintic activity of this extract was evaluated on Indian adult earthworms, *Pheretima posthuma*, and exhibited a dose dependent inhibition of spontaneous motility (paralysis), and evoked responses to pin-prick, and the effects were comparable with that of piperazine citrate. The extract were also assayed for antimicrobial activity against various Gram positive organisms such as *Staphylococcus epidermidis*, *Micrococcus luteus*, *Bacillus subtilis*, and Gram negative organisms such as *Escherichia coli*, *Pseudomonas vulgaris*, *Salmonella typhi*, and fungal strains *Aspergillus niger*, and *Candida albicans*. Antimicrobial activity was conducted by the agar well diffusion method. The extract showed varies levels of antimicrobial activity on different test microorganisms. Future studies are in process to isolate the active principles responsible for the activity.

Keywords: Petroleum ether extract, *Bruguiera gymnorrhiza (L.) Lam.* (Rhizophoraceae), antimicrobial activity, anthelmintic activity, leaves.

INTRODUCTION

Bruguiera gymnorrhiza (L.) Lam. is a plant of the Rhizophoraceae family native along the inland margin of mangrove swamps, and occasionally along beaches. Bark contains D-glucose, rhamnose, arabinose, tannins, a mixture of bruguierol and isobruguierol [1-3]. Apart from that alphaamyrin, beta-amyrin, lupeol, oleanolic acid, ursolic acid, taraxerol, gymnorrhizol. Ellagic acid and derivatives are also present. The leaves have antimicrobial activity(4). The dried wood is insecticidal. The root is used to restore lost appetite and is used to treat diabetes [1]. Prevalence of human helminthiasis in India is as high as 70 to 80 % of which ascariasis is most common, and it exacts a heavy toll on human health and productivity. Although effective anthelmintics are currently available for most infections, the search of novel compounds is essential in view of the development of resistanceas noted in animals [5, 6]. The objective of this work was to explore the anthelmintic and

antimicrobial properties of peels of *Bruguiera gymnorrhiza (L.) Lam.*

MATERIALS AND METHODS

Plant material

The Leaves of *Bruguiera gymnorrhiza (L.) Lam.* were collected in month of January from Tirupati, Chittoor (Dist), Andhra Pradesh, India. The taxonomical identification of the plant was done by Dr. K. Madhava chetty, Assistant Professor, Department of Botany, Sri Venkateshwara University (SVU), Tirupati. The voucher specimen of the plant was deposited at the college for further reference.

Extraction and isolation

Fresh leaves (300 g) were homogenized with 600 mL of petroleum ether for 5 min in a blender. The homogenate was placed in a 2 L Erlenmeyer flask. Three

hundred milliliters of petroleum ether were added in the flask and further extracted three times for 10 min each in an ultrasonic bath at room temperature. The extract was filtered and evaporated under vacuum at 35°C to dryness. The residue (2.22 g) was collected and stored at -20°C until use.

Test microorganism

The microorganisms used for the antimicrobial activity evaluation were obtained from the Microbiology Laboratory, Sree Vidyanikethan College of Pharmacy, Sree Sainath nagar, Tirupati -517-102, Andhra Pradesh, India. They were *Staphylococcus epidermidis* SMC 65, *Micrococcus luteus* MLM 541, *Bacillus subtilis* BSCC 87, *Escherichia coli* ECM 453, *Pseudomonas vulgaris* PVS 01, *Salmonella typhi* TSP 501, *Candida albicans* CAS 22 and *Aspergillus niger* ANG 432.

Antimicrobial activity

The agar diffusion method was used for the antimicrobial activity evaluations (7). Wells of 8 mm diameter were punched into the Mueller-Hinton Agar (MHA, Merck), having the test microorganism and filled with 100 mg/ml of petroleum ether extract. The plates were incubated for 18 h at 37 °C. Antimicrobial activity was evaluated by measuring the inhibition zone (including 8 mm diameter wells) against the test microorganisms. Standard antibiotic discs Ciprofloxacin (25 mg) and Griseofulvin (25 mg) was used as a reference.

Anthelmintic activity

The anthelmintic activity was assessed using adult indian earthworms, *Pheretima posthuma* due to its anatomical and physiological resemblance with the intestinal roundworm parasites of human being (8). The method of Dash *et al.*, (9) was followed for anthelmintic screening. Groups are divided into seven, each group consisting of six earthworms of approximately equal size. Each group was treated with one of the following: vehicle (1% gum acacia in normal saline), piperazine citrate (10, 20, 50 mg/ml) and petroleum ether extract of *Bruguiera gymnorrhiza* (L.) Lam. (10, 20, 50 mg/ml) in normal saline containing 1% gum acacia. Observation was made for the time taken to paralysis and/or death of individual worms up to four hours of test period. Paralysis was said to occur when the worms did not revive even in normal saline. Death was concluded when the

worms lost their motility followed with fading away of their body colour.

Results and Discussion

In the search for compounds with anthelmintic activity, a number of substances have been screened using different species of worms, for example, earthworms, *Ascaris*, *Nippostrongylus* and *heterakis*. Of all these species, earthworms have been widely for the initial evaluation of anthelmintic compounds in vitro because they resemble intestinal worms in their reaction to anthelmintics and are easily available. It has been demonstrated that all anthelmintics are toxic to earthworms and a substance toxic to earthworms is worthy for investigation as an anthelmintic (10). In this study we have evaluated the effect petroleum ether extract of *Bruguiera gymnorrhiza* (L.) Lam. On earthworms. The results obtained are summarized in Table 1. It has been noted that petroleum ether extract showed comparable activity with that of standard piperazine citrate, a drug now widely used as anthelmintic, and in which the activity increased with concentration. Furthermore the antimicrobial activity of petroleum ether extract of *Bruguiera gymnorrhiza* (L.) Lam. was examined and found to exhibit good antibacterial activity against most of the Gram positive and Gram negative organisms which has been depicted in the Table 2. Among the test organisms the extract showed good antimicrobial activity against *Staphylococcus epidermis*, *Micrococcus luteus* and *Pseudomonas vulgaris*, and moderate activity against *Escherichia coli* and *Salmonella typhi*, and no activity against *Bacillus subtilis*. The extract showed good antifungal activity against *Candida albicans* and no activity against *Aspergillus niger*. The result of the antimicrobial activity expressed in terms of diameter of zone of inhibition in millimeter. The performance of *Bruguiera gymnorrhiza* (L.) Lam. extract against sensitive bacteria isolates did not show difference when compared with established commercial antibiotics prepared with amoxicillin and griseofulvin (Table 2).

These results suggests the need for further studies on this extract to identify, isolate, characterize and elucidate the structure of the active ingredient(s) using some spectroscopic techniques such as infrared spectrometry, nuclear magnetic resonance spectroscopy and mass spectrometry.

Table 1. Anthelmintic activity of petroleum ether extract of *Bruguiera gymnorrhiza* (L.) Lam.

| Treatment | Time taken for paralysis (min) | Time taken for death (min) |
|-----------------------------|--------------------------------|----------------------------|
| Vehicle | - | - |
| Petroleum ether extract | 90.60 ± 1.23 | 218.83 ± 2.06 |
| 20 mg/ml | 54.17 ± 1.40 | 183.14 ± 1.64 |
| 50mg/ml | 22.63 ± 1.45 | 56.53 ± 1.45 |
| Piperazine citrate 10 mg/ml | 101.0 ± 1.14 | - |
| 20 mg/ml | 74.38 ± 1.46 | - |
| 50mg/ml | 36.24 ± 0.49 | - |

Values represent the mean ± SD from six observations

Table 2. Antimicrobial Properties of Leaves of *Bruguiera gymnorrhiza* (L.) Lam.

| Organisms | Mean zone of inhibition (mm) | | | |
|-----------------------------------|------------------------------|---------|-----------------------|----------------------|
| | Petroleum extract | Control | Ciprofloxacin (25 µg) | Griseofulvin (25 µg) |
| <i>Staphylococcus epidermidis</i> | 28 | - | 32 | NT |
| <i>Micrococcus luteus</i> | 22 | - | 26 | NT |
| <i>Bacillus subtilis</i> | - | - | 29 | NT |
| <i>Escherichia coli</i> | 16 | - | 31 | NT |
| <i>Pseudomonas vulgaris</i> | 29 | - | 33 | NT |
| <i>Salmonella typhi</i> | 18 | - | 29 | NT |
| <i>Candida albicans</i> | 16 | - | NT | 17 |
| <i>Aspergillus niger</i> | - | - | NT | 12 |

NT – not tested

REFERENCES

1. Cambie RC. & Ash J. Fijian Medicinal Plants, CSIRO, Australia, 1994, 102-103.
2. Subba Rao GSR. *et al.*, *Indian J. Chem. Sec. B*, 25 (2), 1986, 113-122.
3. Locher CP. *et al.*, *Phytomedicine*, 2, 1996, 259-264.
4. Whistler WA. *J. Ethnopharmacol*, 13 (3), 1985, 239-280.
5. Prichard R. Anthelmintic resistance, *Vet Parasitol* 54, 1994, 259-268.
6. Al-Qarawi AA, Mahmoud OM, Sobaih Haroun EM, Adam SE. A preliminary study on the activity of *Calotropis procera* latex against *Haemonchus contortus* infection in najdi sheep. *Vet Res Commun*, 25, 2001, 131-134.
7. Maruzzella JC, Henry PA. The antimicrobial activity of perfume oils. *Journal of the American Pharmaceutical Association*, 28(7), 1958, 471.
8. Thorn GW, Adams RD, Braunwald E, Isselbacher KJ, Petersdorf RG. *Harrisons principle of internal medicine*, Mcgraw Hill Co, New York, 1977, 1088.
9. Dash GK, Mishra B, Panda A, Patro CP, Ganapaty S. *Indian J Nat Products* 19(3), 2003, 24-26.
10. Sollmann T. Anthelmintics and their efficiency as tested on earthworms. *J Pharmacol Exp Ther*, 12, 1918, 129-170.